

Non-specific urethritis (NSU) and Cervicitis

NSU is the term used to describe swelling or inflammation of the urethra (the tube inside the penis) after other specific infections have been tested for and given a negative result, such as chlamydia and gonorrhoea. NSU is not very common, as such infections are usually easy to detect.

Cervicitis ((inflammation of the cervix, or neck of the womb) is the female equivalent of NSU.

Causes

NSU and Cervicitis can be caused by a number of different organisms, including sexually transmissible infections (STIs).

Symptoms

Symptoms can include irritation when passing urine or discharge from the penis or vagina. Females may also experience lower abdominal pain or pain during sex. Symptoms are similar to those of many STIs, and most of the time they indicate chlamydia or gonorrhoea.

How is it transmitted?

NSU and Cervicitis is transmitted through an exchange of body fluids (semen, blood and vaginal fluids) during unprotected vaginal, anal or oral sexual activity.

How do I know if I have it?

Testing for NSU and Cervicitis involves a swab from the cervix and/or urethra. **For more information about STI testing go to www.fpwa.org.au/stitests.htm**

Treatment/Management

Treatment for NSU and Cervicitis involves a course of antibiotics. All sexual partners should also be tested and treated. Sex should be avoided until the treatment is complete.

Why is treatment important?

If left untreated, NSU and Cervicitis can lead to infertility in both men and women. In males it can also lead to prostatitis (inflammation of the prostate gland) and epididymitis (inflammation of the testicles).

Prevention

Using condoms and dams reduces the risk of contracting NSU or Cervicitis (a dam is a thin latex square held over the vaginal or anal area during oral sex). They stop body fluids like semen, blood and vaginal fluids from being exchanged. **For more information about condoms and dams go to www.fpwa.org.au/healthinformation/informationsheets/**

**For more information about NSU or Cervicitis contact the Sexual Health Helpline on
9227 6178 or 1800 198 205 (country callers) or email sexhelp@fpwa.org.au**

**Practising safe sex reduces the risk of contracting HIV and other
sexually transmissible infections (STIs).**



70 Roe Street, PO Box 141, Northbridge, WA, 6865
ph: (08) 9227 6177 fax: (08) 9227 6871
www.fpwa.org.au

info@fpwa.org.au

Sexual Health Helpline (08) 92276178 or 1800 198 205 (Country Callers)
sexhelp@fpwa.org.au

Quarry Health Centre for under 25s
7 Quarry Street, PO Box 378, Fremantle, WA, 6959
ph: (08) 9430 4544 fax: (08) 9430 4544
quarry@fpwa.org.au

© Family Planning Association of WA (Inc) March 2006